

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB05706 - Goat Anti-GCIP / MAID Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CCNDBP1, cyclin D-type binding-protein 1, DIP1, GCIP, HHM Protein, MAID protein, grap2 cyclin interacting protein, D-type cyclin-interacting protein 1

Official Symbol: CCNDBP1

Accession Number(s): NP_036274.3; NP_411241.1

Human GeneID(s): 23582

Important Comments: This antibody is expected to recognise both human isoforms of

this protein.

Immunogen

Peptide with sequence C-NRIKELTQSELEL, from the C Terminus of the protein sequence according to NP_036274.3; NP_411241.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Preliminary experiments gave an approx 55kDa band in K562 lysate at 0.5 ug/ml. This band was successfully blocked by incubation with the immunising peptide. Please note that currently we cannot find an explanation in the literature for the band we observe given the predicted size of approx. 41kDda and 27kDa according to NP_036274; NP_411241. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

IHC: In paraffin embedded Human Prostate shows textured cytoplasm staining with emphasis on the luminal side of the secretory cells in the glands. Recommended concentration, 3-6µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Dog



 $EB05706~(3.8\mu g/ml)~staining~of~paraffin~embedded~Human~Prostate.~Steamed~antigen~retrieval~with~citrate~buffer~pH~6,~AP-staining.$