



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB07027 - Goat Anti-GLI1 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** GLI1, glioma-associated oncogene homolog 1 (zinc finger protein), HGNC:4317, GLI, glioma-associated oncogene homolog (zinc finger protein), glioma-associated oncogene homolog 1, Tax Helper Protein, GLI family zinc finger 1, glioma-associated oncogene family zinc finger 1, glioma-associated oncogene homolog 1 (zinc finger protein)

**Official Symbol:** GLI1

**Accession Number(s):** NP\_005260.1; NP\_001153517.1; NP\_001161081.1

**Human GeneID(s):** [2735](#)

### Immunogen

Peptide with sequence C-EDAPAQEPSYQSPK, from the internal region of the protein sequence according to NP\_005260.1; NP\_001153517.1; NP\_001161081.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:8000.

**Immunofluorescence:** Strong expression of the protein seen in the cytoplasm and nuclei of U2OS cells. Recommended concentration: 10µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB07027 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).