

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07027 - Goat Anti-GLI1 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** GLI1, glioma-associated oncogene homolog 1 (zinc finger protein), HGNC:4317, GLI, glioma-associated oncogene homolog (zinc finger protein), glioma-associated oncogene homolog 1, Tax Helper Protein, GLI family zinc finger 1, glioma-associated oncogene family zinc finger 1, glioma-associated oncogene homolog 1 (zinc finger protein)

Official Symbol: GLI1

Accession Number(s): NP\_005260.1; NP\_001153517.1; NP\_001161081.1

Human GeneID(s): 2735

## Immunogen

Peptide with sequence C-EDAPAQEPSYQSPK, from the internal region of the protein sequence according to NP\_005260.1; NP\_001153517.1; NP\_001161081.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:8000.

**Immunofluorescence:** Strong expression of the protein seen in the cytoplasm and nuclei of U2OS cells. Recommended concentration:  $10\mu g/ml$ .

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB07027 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).