

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB10742 - Goat Anti-GNAQ / ALPHA-q (aa162-175) Antibody

Size: 100µg specific antibody in 200µl

Target Protein

Principal Names: GNAQ, guanine nucleotide binding protein (G protein), q polypeptide,

G-ALPHA-q, GAQ
Official Symbol: GNAQ

Accession Number(s): NP_002063.2

Human GeneID(s): 2776

Non-Human GenelD(s): 14682 (mouse), 81666 (rat)

Immunogen

Peptide with sequence C-NDLDRVADPAYLPT, from the internal region of the protein sequence according to NP_002063.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 40kDa band observed in Human, Mouse and Rat Testis lysates, in Pig Colon and Spleen lysates and in lysates of Human Peripheral Blood Mononucleocytes (calculated MW of 42.1kDa according to Human NP_002063.2 and to Rat NP_112298.1).


Recommended concentration: 0.1-0.3µg/ml.

IHC: Paraffin embedded Human Kidney. Recommended concentration: 5µg/ml.


Species Reactivity


Tested: Human, Mouse, Rat, Pig

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow




EB10742 (0.1μg/ml) staining of Mouse (A) and Rat (B) Testis lysates (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.





EB10742 (0.1μg/ml) staining of Pig Colon (A) and Pig Spleen (B) lysates (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB10742 (5µg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.