

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB12564 - Goat Anti-GNL3 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** GNL3, guanine nucleotide binding protein-like 3 (nucleolar), C77032, E2IG3, NNP47, NS, E2-induced gene 3 protein, estradiol-induced nucleotide binding protein, guanine nucleotide-binding protein-like 3, novel nucleolar protein 47, nucleolar GTP-binding protein 3, nucleostemin

Official Symbol: GNL3

Accession Number(s): NP\_055181.3; NP\_996561.1

Human GeneID(s): 26354

Important Comments: This antibody is expected to recognize both reported isoforms

(NP\_055181.3; NP\_996561.1). Reported variants represent identical protein:

NP\_996561.1, NP\_996562.1.

## **Immunogen**

Peptide with sequence C-EKNNAQSIRAIKGPH, from the internal region of the protein sequence according to NP\_055181.3; NP\_996561.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot**: Approx 65+75kDa bands observed in nuclear lysates of cell lines HeLa, HepG2 and K562 (calculated MW of 62.0kDa according to NP\_055181.3). The observed 75kDa band likely represents the acetylated form. Recommended concentration: 1-3µg/ml.

# **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB12564 (2μg/ml) staining of HeLa (A)HepG2 (B) and K562 (C) nuclear lysates (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.