



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06926 - Goat Anti-GOLGA3 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: GOLGA3, HGNC:4426, GCP170, MEA-2, Golgi autoantigen, golgin subfamily a, 3, Golgi complex-associated protein of 170 kD, Golgi membrane associated protein, Golgi peripheral membrane protein, SY2/SY10 protein, golgin-160, golgin-165, male enhanced antigen-2

Official Symbol: GOLGA3

Accession Number(s): NP_005886.2

Human GeneID(s): [2802](#)

Immunogen

Peptide with sequence C-HSQSRASKEGPGE, from the C Terminus of the protein sequence according to NP_005886.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 170kDa band observed in human HeLa lysates (calculated MW of 167kDa according to NP_005886.2). Recommended concentration: 0.1-0.3µg/ml. An additional band of unknown identity was also consistently observed at approx 25kDa. This band was successfully blocked by incubation with the immunising peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB06926 (0.1µg/ml) staining of Hela lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.