



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB07290 - Goat Anti-GPR94 / TRA1 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** TRA1, GRP94, tumor rejection antigen (gp96) 1, HGNC:12028, ECGP, GP96, Tumor rejection antigen-1 (gp96), endothelial cell (HBMEC) glycoprotein, glucose regulated protein, 94 kDa, tumor rejection antigen (gp96) 1

**Official Symbol:** HSP90B1

**Accession Number(s):** NP\_003290.1

**Human GeneID(s):** [7184](#)

**Non-Human GeneID(s):** 22027 (mouse), 362862 (rat)

### Immunogen

Peptide with sequence C-KEGVKFDSEKTK, from the internal region of the protein sequence according to NP\_003290.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:8000.

**Western blot:** Approx. 100kDa band observed in lysates of cell lines HeLa, HepG2 and U2OS and in preliminary testing of NIH3T3 cell lysate and Mouse Liver lysate (calculated MW of 92.5kDa according to Human NP\_003290.1). This molecular weight is routinely observed by other sources. Recommended concentration: 0.3-0.5µg/ml. Primary incubation 1 hour at room temperature. **Positive Control:** A batch specific positive control lysate is available for this product. Please contact [Sales@everestbiotech.com](mailto:Sales@everestbiotech.com) for availability.

**IHC:** Paraffin embedded Human Liver. Recommended concentration: 5-7µg/ml.

**Flow Cytometry:** Flow cytometric analysis of HeLa cells. Recommended concentration: 10ug/ml.


### Species Reactivity

**Tested:** Human


**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Pig, Cow, Zebrafish




EB07290 (0.5 $\mu$ g/ml) staining of HeLa (A) and U2OS (B), and (0.3 $\mu$ g/ml) on HepG2 (C), cell lysate (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.



EB07290 (6 $\mu$ g/ml) staining of paraffin embedded Human Liver. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB07290 Negative Control showing staining of paraffin embedded Human Liver, with no primary antibody.



EB07290 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.