



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12813 - Goat Anti-HAO2 (aa54-67) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: HAO2, hydroxyacid oxidase 2 (long chain), GIG16, HAOX2, (S)-2-hydroxy-acid oxidase, peroxisomal, cell growth-inhibiting gene 16 protein, glycolate oxidase, hydroxyacid oxidase 2, long chain alpha-hydroxy acid oxidase, long-chain L-2-hydroxy acid oxidase

Official Symbol: HAO2

Accession Number(s): NP_057611.1

Human GeneID(s): [51179](#)

Important Comments: Reported variants represent identical protein: NP_001005783.1, NP_057611.1.

Immunogen

Peptide with sequence C-DVSEVDTRTTIQGE, from the internal region of the protein sequence according to NP_057611.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 38kDa band observed in Human Kidney lysates (calculated MW of 38.8kDa according to NP_057611.1). Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB12813 (0.3 μ g/ml) staining of Human Kidney lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.