



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05610 - Goat Anti-HEC1 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** NDC80, HEC1, HEC, highly expressed in cancer, rich in leucine heptad repeats, HEC-1, KNTC2, NDC80 homolog, kinetochore complex component (S. cerevisiae), TID3, hsNDC80, kinetochore associated 2

**Official Symbol:** NDC80

**Accession Number(s):** NP\_006092.1

**Human GeneID(s):** [10403](#)

### Immunogen

Peptide with sequence C-YEKKATLIKSSEE, from the C Terminus of the protein sequence according to NP\_006092.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.


**Western blot:** Approx 75-80kDa band observed in Hela lysate (predicted MW of 74kDa according to NP\_006092). Recommended for use at 0.5-2µg/ml.

**IHC:** Paraffin embedded Human Liver and Adrenal Gland. Recommended concentration: 5µg/ml.


### Species Reactivity

**Tested:** Human


**Expected from sequence similarity:** Human



EB05610 staining (0.5µg/ml) of Hela lysate (RIPA buffer, 35µg total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.



EB05610 (5µg/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB05610 (5µg/ml) staining of paraffin embedded Human Adrenal Gland. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.