



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06609 - Goat Anti-HIP14 / ZDHHC17 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** ZDHHC17, HIP14, HIP3, HYPH, KIAA0946, zinc finger, DHHC domain containing 17, Huntingtin interacting protein H, huntingtin interacting protein 3, huntingtin interacting protein 14, HSPC294, zinc finger, DHHC-type containing 17

**Official Symbol:** ZDHHC17

**Accession Number(s):** NP\_056151.2

**Human GeneID(s):** [23390](#)

### Immunogen

Peptide with sequence CYDQISGSGYQLV, from the C Terminus of the protein sequence according to NP\_056151.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:8000.

**Western blot:** Approx 70kDa band observed in Mouse Brain lysates (calculated MW of 72.7kDa according to NP\_056151.1). Recommended concentration: 1-3µg/ml.

**IHC:** Paraffin embedded Human Testis. Recommended concentration: 5µg/ml.


### Species Reactivity

**Tested:** Human, Mouse

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow



EB06609 (1 $\mu$ g/ml) staining of Mouse Brain lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB06609 (5 $\mu$ g/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.