

#### **International Office**

Everest Biotech Ltd Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service: <u>customerservice@vectorlabs.com</u> Technical Service: <u>technical@vectorlabs.com</u>

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06640 - Goat Anti-HIP14L / ZDHHC13 Antibody

Size: 100µg specific antibody in 200µl

# **C**

## **Target Protein**

Principal Names: ZDHHC13, HIP14L, HIP3RP, FLJ10852, FLJ10941, MGC64994, zinc finger, DHHC domain containing 13, HIP14-like, HIP14-related protein, huntingtin interacting protein HIP3RP, hypothetical protein similar to ankyrin repeat-containing priotein AKR1, zinc finger, DHHC-type containing 13
Official Symbol: ZDHHC13
Accession Number(s): NP\_061901.2; NP\_001001483.1
Human GenelD(s): 54503
Non-Human GenelD(s): 243983 (mouse), 365252 (rat)
Important Comments: This antibody is expected to recognise both reported isoforms (NP\_061901.2 and NP\_001001483.1).

### Immunogen

Peptide with sequence C-FHPAREKVLRSV, from the C Terminus of the protein sequence according to NP\_061901.2; NP\_001001483.1.

Please note the peptide is available for sale.

#### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx. 70kDa band observed in Human Brain (Cerebellum) lysates (calculated MW of 70.9kDa according to NP\_061901.2). Recommended concentration: 1-3µg/ml.


IHC: Paraffin embedded Human Lung. Recommended concentration:  $5\mu$ g/ml.

### **Species Reactivity**

Tested: Human Expected from sequence similarity: Human, Mouse, Rat



EB06640 (0.1µg/ml) staining of Human Cerebellum lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB06640 (5µg/ml) staining of paraffin embedded Human Lung. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.