



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06904 - Goat Anti-Histamine Receptor H1 (C Term) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: HRH1, histamine receptor H1, HGNC:5182, H1-R, hisH1, histamine receptor, subclass H1, OTTHUMP00000160133, histamine H(1) receptor

Official Symbol: HRH1

Accession Number(s): NP_000852.1; NP_001091681.1; NP_001091682.1; NP_001091683.1

Human GeneID(s): [3269](#)

Non-Human GeneID(s): 15465 (mouse), 24448 (rat)

Important Comments: Variants (NP_000852.1; NP_001091681.1; NP_001091682.1; NP_001091683.1) encode the same protein.

Immunogen

Peptide with sequence CNENFKKTFKRILH, from the C Terminus of the protein sequence according to NP_000852.1; NP_001091681.1; NP_001091682.1; NP_001091683.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

IHC: This product has been successfully used in IHC on Rat (PMID: 30226827).

Immunofluorescence: This product has been successfully used in IF on Rat (PMID: 30413645 and 30143981).

Flow Cytometry: Flow cytometric analysis of A431 cells. Recommended concentration: 10ug/ml.

Species Reactivity

Tested: Human, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

Specific References

This antibody has been successfully used in IF on Rat:

Chen ZP, Zhang XY, Peng SY, Yang ZQ, Wang YB, Zhang YX, Chen X, Wang JJ, Zhu JN
Histamine H1 Receptor Contributes to Vestibular Compensation.

J Neurosci. 2019 Jan 16;39(3):420-433.

PMID: 30413645

This antibody has been successfully used in IHC on Rat:

Zhuang QX, Li GY, Li B, Zhang CZ, Zhang XY, Xi K, Li HZ, Wang JJ, Zhu JN.

Regularizing firing patterns of rat subthalamic neurons ameliorates parkinsonian motor deficits.

J Clin Invest. 2018 Sep 18.

PMID: 30226827


This antibody has been successfully used in IF on Rat:

Miao-Jin Ji, Xiao-Yang Zhang, Xiao-Chun Peng, Yang-Xun Zhang, Zi Chen, Lei Yu,
Jian-Jun Wang, Jing-Ning Zhu

Histamine Excites Rat GABAergic Ventral Pallidum Neurons via Co-activation of H1 and
H2 Receptors

Neurosci Bull. 2018 Dec;34(6):1029-1036.

PMID: 30143981



EB06904 Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.