

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB12827 - Goat Anti-HLA-B Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** HLA-B, major histocompatibility complex, class I, B, AS, HLAB, SPDA1, HLA class I histocompatibility antigen, B alpha chain, MHC Class I HLA heavy chain, MHC HLA-B cell surface glycoprotein, MHC HLA-B transmembrane glycoprotein,

MHC class I antigen GN0010 **Official Symbol:** HLA-B

Accession Number(s): NP\_005505.2

Human GeneID(s): 3106

### **Immunogen**

Peptide with sequence C-DRNTQIYKAQAQT, from the internal region of the protein sequence according to NP\_005505.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:64000.

**Western blot:** Approx 40+45kDa bands observed in lysates of cell line Jurkat (calculated MW of 40.5kDa according to NP\_005505.2). The observed 45kDa band represents the glycosylated form. Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Tonsil. Recommended concentration: 5µg/ml.


### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



 $EB12827\ (0.1\mu\text{g/ml})\ staining\ of\ Jurkat\ lysate\ (35\mu\text{g}\ protein\ in\ RIPA\ buffer).\ Detected\ by\ chemiluminescence.$ 



EB12827 (5μg/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.