

#### International Office

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09567 - Goat Anti-HMGA1 (aa9-21) Antibody

Size: 100µg specific antibody in 200µl



#### **Target Protein**

**Principal Names:** HMGA1, high mobility group AT-hook 1, HMG-R, HMGA1A, HMGIY, MGC12816, MGC4242, MGC4854, OTTHUMP00000016222, OTTHUMP00000016223, OTTHUMP00000016224, OTTHUMP00000039618, high-mobility group (nonhistone chromosomal) protein isoforms I and Y, nonhistone chromosomal high-mobility group

protein HMG-I/HMG-Y
Official Symbol: HMGA1

Accession Number(s): NP\_665906.1; NP\_002122.1

Human GenelD(s): 3159

Non-Human GenelD(s): 15361 (mouse), 117062 (rat)

**Important Comments:** This antibody is expected to recognize reported isoforms a (NP\_665906.1, NP\_665908.1, NP\_665911.1) and b (NP\_002122.1, NP\_665910.1, NP\_665910.1).

NP\_665912.1). This antibody is not expected to cross-react with HMGA2.

### **Immunogen**

Peptide with sequence C-SQPLASKQEKDGT, from the internal region (near N Terminus) of the protein sequence according to NP\_665906.1; NP\_002122.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.


Western blot: Approx 24kDa band observed in Human Pancreas (calculated MW of 11.7kDa/10.7kDa according to NP\_665906.1; NP\_002122.1). This molecular weight is routinely observed by other sources. This band was successfully blocked by incubation with the immunizing peptide. Recommended concentration: 0.5-1μg/ml. Primary incubation was 1 hour.

**IHC:** Paraffin embedded Human Skin and Placenta. Recommended concentration: 3.75µg/ml.


## **Species Reactivity**

Tested: Human


Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow, Pig



EB09567 (0.5 $\mu$ g/ml) staining of Human Pancreas lysate (35 $\mu$ g protein in RIPA buffer) with (B) and without (A) blocking with the immunising peptide. Detected by chemiluminescence.



EB09567 (3.75 $\mu$ g/ml) staining of paraffin embedded Human Skin. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB09567 (3.75 $\mu$ g/ml) staining of paraffin embedded Human Placenta. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.