

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB12202 - Goat Anti-HOXB6 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** HOXB6, homeobox B6, HOX2, HOX2B, HU-2, Hox-2.2, homeo box 2B, homeo box B6, homeobox protein Hox-2.2, homeobox protein Hox-2B, homeobox

protein Hox-B6, homeobox protein Hu-2

Official Symbol: HOXB6

Accession Number(s): NP\_061825.2

Human GeneID(s): 3216

Non-Human GenelD(s): 15414 (mouse), 497986 (rat)

## **Immunogen**

Peptide with sequence EPRKSDCAQDKS, from the internal region of the protein sequence according to NP\_061825.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:1000.

**Western blot:** Approx 26kDa band observed in lysates of cell lines HEK293, CACO-2 and MRC5 PD19 (calculated MW of 25.4kDa according to NP\_061825.2). Recommended

concentration: 1-3µg/ml.


IHC: In paraffin embedded Human Lung shows nuclear staining in the alvioli.

Recommended concentration, 5-10µg/ml.

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB12202 (5µg/ml) staining of paraffin embedded Human Lung. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB12202 (1 $\mu$ g/ml) staining of HEK293 lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.