



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09834 - Goat Anti-HOXD13 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: BDE, BDSD, homeo box 4l, homeo box D13, homeobox D13, HOX4l, SPD, HOXD13

Official Symbol: HOXD13

Accession Number(s): NP_000514.2

Human GeneID(s): [3239](#)

Non-Human GeneID(s): 15433 (mouse), 288154 (rat)

Immunogen

Peptide with sequence C-HGVGLQQNALKSSPH, from the internal region of the protein sequence according to NP_000514.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 37kDa band observed in Rat Brain lysates (calculated MW of 34.9kDa according to NP_001099356.1). Recommended concentration: 0.3-1µg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Prostate. Recommended concentration: 8µg/ml.

Immunofluorescence: Strong expression of the protein seen in the nucleus of HEK293 and U2OS cells. Recommended concentration: 10µg/ml.


Species Reactivity

Tested: Human, Rat


Expected from sequence similarity: Human, Mouse, Rat, Cow, Pig




EB09834 (1 μ g/ml) staining of Rat Brain lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.




EB09834 Immunofluorescence analysis of paraformaldehyde fixed HEK293 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (2 μ g/ml), showing nuclear staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (2 μ g/ml).



EB09834 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing strong nuclear and weak cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB09834 (8µg/ml) staining of paraffin embedded Human Prostate. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB09834 Negative Control showing staining of paraffin embedded Human Prostate, with no primary antibody.