

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB11196 - Goat Anti-HP beta chain Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** binding peptide, BP, haptoglobin, haptoglobin alpha(1S)-beta, haptoglobin alpha(2FS)-beta, haptoglobin, alpha polypeptide, haptoglobin, beta

polypeptide, HP2ALPHA2, HPA1S, MGC111141, HP

Official Symbol: HP

Accession Number(s): NP\_005134.1; NP\_001119574.1

Human GeneID(s): 3240

Important Comments: This antibody is expected to recognize both reported isoforms

(NP\_005134.1; NP\_001119574.1).

## Immunogen

Peptide with sequence C-STVPEKKTPKSP, from the internal region of the protein sequence according to NP\_005134.1; NP\_001119574.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:8000.

**Western blot:** Approx 37+40kDa bands observed in Human Liver lysates and 40kDa in Pig Liver lysates (calculated MW of 45.2kDa according to NP\_005134.1 and 38.5kDa according to NP\_001119574.1). Recommended concentration: 0.1-0.3μg/ml.


#### **Species Reactivity**

Tested: Human, Pig

Expected from sequence similarity: Human, Dog, Pig



EB11196 (0.1 $\mu$ g/ml) staining of Human Liver lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB11196 (1µg/ml) staining of Pig Liver lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.