

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB11978 - Goat Anti-HSPA13 / STCH (aa83-95) Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** HSPA13, heat shock protein 70kDa family, member 13, STCH, heat shock 70 kDa protein 13, microsomal stress 70 protein ATPase core, microsomal stress-70 protein ATPase core, stress 70 protein chaperone, microsome-associated, 60kD, stress 70 protein chaperone, microsome-associated, 60kDa, stress-70 protein chaperone microsome-associated 60 kDa protein

Official Symbol: HSPA13

Accession Number(s): NP\_008879.3

Human GeneID(s): 6782

## Immunogen

Peptide with sequence C-ESVELADSNPQNT, from the internal region of the protein sequence according to NP\_008879.3.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot:** Approx 60-65kDa band observed in Human Uterus and Colon lysates (calculated MW of 51.9kDa according to NP\_008879.3)..This band was successfully blocked by incubation with the immunizing peptide and corresponds to different antibodies from other sources. Recommended concentration: 0.1-0.3μg/ml. Primary incubation 1 hour at room temperature.

#### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB11978 (0.1μg/ml) staining of Human Uterus lysate (A) + peptide (B) and (0.3ug/ml) Colon lysate (C) + peptide (D) (35μg protein in RIPA buffer). Detected by chemiluminescence.