

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05765 - Goat Anti-HSPC150 / UBE2T Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** UBE2T, HSPC150, PIG50, HSPC150 protein similar to ubiquitin-conjugating enzyme, ubiquitin conjugating enzyme, ubiquitin-conjugating enzyme E2T (putative), ubiquitin-conjugating enzyme E2T

**Official Symbol:** UBE2T

**Accession Number(s):** NP\_054895.1; NP\_001297255.1

**Human GeneID(s):** [29089](#)

### Immunogen

Peptide with sequence C-QLVGIEKKFHPDV, from the C Terminus of the protein sequence according to NP\_054895.1; NP\_001297255.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx. 23-24kDa band observed in nuclear lysates of cell lines HeLa, Jurkat, HEK293, HepG2 and in THP-1 cell lysates, and approx. 25kDa in nuclear lysates of cell line A431 and in K562, cell lysates (calculated MW of 22.5kDa according to NP\_054895.1). Recommended concentration: 0.1-0.3µg/ml. Primary incubation 1 hour at room temperature.


**Immunofluorescence:** Strong expression of the protein seen in the nuclei of HeLa and U2OS cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of HeLa cells. Recommended concentration: 10µg/ml.

### Species Reactivity


**Tested:** Human

**Expected from sequence similarity:** Human




EB05765 optimised QC. Primary incubation 1 hour at room temperature.

Image A: K562 cell lysate at primary Ab concentration 0.1µg/ml, Image B: THP-1 cell lysate at primary Ab concentration 0.3µg/ml. (Loaded 35µg protein in RIPA buffer, per lane). Detected by chemiluminescence.




EB05765 optimised QC. Primary incubation 1 hour at room temperature.


Images A, B, C, D: A431, HEK293, HeLa, Jurkat nuclear cell lysate at primary Ab concentration 0.3µg/ml, Image E: HepG2 nuclear cell lysate at primary Ab concentration 0.1µg/ml. (Loaded 35µg protein in RIPA buffer, per lane). Detected by chemiluminescence.



EB05765 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing strong nuclear staining. Actin filaments were stained with phalloidin (red) and the nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB05765 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing strong nuclear staining. Actin filaments were stained with phalloidin (red) and the nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB05765 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.