

#### **UK Office**

### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07999 - Goat Anti-HTATSF1 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** HTATSF1, HIV-1 Tat specific factor 1, RP1-196E23.2, TAT-SF1, dJ196E23.2, HIV TAT specific factor 1, cofactor required for Tat activation of HIV-1

transcription

Official Symbol: HTATSF1

Accession Number(s): NP\_055315.2

Human GeneID(s): 27336

Non-Human GenelD(s): 317612 (rat)

# **Immunogen**

Peptide with sequence C-QELYGDGKDGDTQTD, from the internal region (near the N Terminus) of the protein sequence according to NP\_055315.2.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 140kDa band observed in lysates of cell line Daudi (calculated MW of 85.9kDa according to NP\_055315.2). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Zhou and Sharp, Science. 1996 Oct 25;274(5287):605-10.; PMID: 8849451). Recommended concentration: 1-3µg/ml. An additional band of unknown identity was also consistently observed at 20kDa. This band was successfully blocked by incubation with the immunising peptide.

**Additional validation:** This antibody has been successfully used in the following paper: Sikorski et al. (2018) PMID: 30377371.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Rat, Dog

### Specific Reference


## This antibody has been successfully used in the following paper:

Krzysztof Sikorski, Adi Mehta, Marit Inngjerdingen, Flourina Thakor, Simon Kling, Tomas Kalina, Tuula A. Nyman, Maria Ekman Stensland, Wei Zhou, Gustavo A. De Souza, Lars Holden, Jan Stuchly, Markus Templin and Fridtjof Lund-Johansen

A high-throughput pipeline for validation of antibodies

Nat Methods. 2018 Nov;15(11):909-912

PMID: 30377371



EB07999 (1 $\mu$ g/ml) staining of Daudi cell lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.