

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07020 - Goat Anti-HVEM / TR2 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** HVEM, TR2, TNFRSF14, tumor necrosis factor receptor superfamily, member 14 (herpesvirus entry mediator), HGNC:11912, ATAR, HVEA, LIGHTR, CD40-like protein precursor, herpesvirus entry mediator, herpesvirus entry mediator A, tumor necrosis factor receptor superfamily, member 14, tumor necrosis factor receptor-like gene2

Official Symbol: TNFRSF14

Accession Number(s): NP\_003811.2

Human GeneID(s): 8764

### Immunogen

Peptide with sequence C-SVQRKRQEAEGE, from the internal region of the protein sequence according to NP\_003811.2.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 30kDa band observed in Human Thymus and Tonsil lysates (calculated MW of 30.4kDa according to NP\_003811.2). Recommended concentration:

 $0.03-0.1 \mu g/ml$ .

IHC: Paraffin embedded Human Kidney. Recommended concentration: 5µg/ml.


### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB07020 (0.03μg/ml) staining of Human Thymus lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB07020 (5μg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.