

### **UK Office**

### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

#### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08424 - Goat Anti-IA2 Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** IA-2, protein tyrosine phosphatase, receptor type, N, FLJ16131, IA-2/PTP, IA2, ICA512, R-PTP-N, islet antigen 2, islet cell antigen 512, islet cell

autoantigen 3, protein tyrosine phosphatase-like N

Official Symbol: PTPRN

Accession Number(s): NP\_002837.1

Human GeneID(s): 5798

Non-Human GenelD(s): 19275 (mouse), 116660 (rat)

### **Immunogen**

Peptide with sequence C-QGEGNIKKNRHPD, from the internal region of the protein sequence according to NP\_002837.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 110kDa band observed in lysates of cell line NIH 3T3 (calculated

MW of 106kDa according to Human NP\_002837.1 and Mouse NP\_033011.2).

Recommended concentration: 0.5-1.5µg/ml.

### **Species Reactivity**

Tested: Mouse

Expected from sequence similarity: Human, Mouse, Rat

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB08424 ( $0.5\mu g/ml$ ) staining of NIH 3T3 lysate ( $35\mu g$  protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.