

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB05943 - Goat Anti-ICBP90 / UHRF1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: UHRF1, ubiquitin-like, containing PHD and RING finger domains, 1, Np95, ICBP90, FLJ21925, transcription factor ICBP90, RNF106, MGC138707, RING finger protein 106, inverted CCAAT box-binding protein of 90 kDa, nuclear zinc finger protein Np95, transcription factor ICBP90, ubiquitin-like with PHD and ring finger domains 1, hNP95, E3 ubiquitin-protein ligase UHRF1

Official Symbol: UHRF1

Accession Number(s): NP_001041666.1; NP_037414.3

Human GeneID(s): 29128

Non-Human GenelD(s): 18140 (mouse)

Important Comments: This antibody is expected to recognise both reported isoforms.

Immunogen

Peptide with sequence C-LNQLFPGYGNGR, from the C Terminus of the protein sequence according to NP_001041666.1; NP_037414.3.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:2000.

Western blot: Approx 100kDa band observed in nuclear lysates of cell line Jurkat (calculated MW of 91.1kDa according to NP_037414.3). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Hopfner, Cancer Res. 2000 Jan 1;60(1):121-8. PMID: 10646863). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa

25kDa

20kDa

15kDa

EB05943 (2μg/ml) staining of nuclear Jurkat lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour.

Detected by chemiluminescence.