

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05943 - Goat Anti-ICBP90 / UHRF1 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** UHRF1, ubiquitin-like, containing PHD and RING finger domains, 1, Np95, ICBP90, FLJ21925, transcription factor ICBP90, RNF106, MGC138707, RING finger protein 106, inverted CCAAT box-binding protein of 90 kDa, nuclear zinc finger protein Np95, transcription factor ICBP90, ubiquitin-like with PHD and ring finger domains 1, hNP95, E3 ubiquitin-protein ligase UHRF1

**Official Symbol:** UHRF1

**Accession Number(s):** NP\_001041666.1; NP\_037414.3

**Human GeneID(s):** [29128](#)

**Non-Human GeneID(s):** 18140 (mouse)

**Important Comments:** This antibody is expected to recognise both reported isoforms.

### Immunogen

Peptide with sequence C-LNQLFPGYGNGR, from the C Terminus of the protein sequence according to NP\_001041666.1; NP\_037414.3.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:2000.

**Western blot:** Approx 100kDa band observed in nuclear lysates of cell line Jurkat (calculated MW of 91.1kDa according to NP\_037414.3). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Hopfner, Cancer Res. 2000 Jan 1;60(1):121-8. PMID: 10646863). Recommended concentration: 1-3µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog



EB05943 (2 $\mu$ g/ml) staining of nuclear Jurkat lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour.  
Detected by chemiluminescence.