



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09484 - Goat Anti-ICOS Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ICOS, inducible T-cell co-stimulator, AILIM, CD278, MGC39850, activation-inducible lymphocyte immunomediatory molecule, inducible costimulator

Official Symbol: ICOS

Accession Number(s): NP_036224.1

Human GeneID(s): [29851](#)

Non-Human GeneID(s): 54167 (mouse)

Immunogen

Peptide with sequence KGGQILCDLTKTKGS, from the internal region of the protein sequence according to NP_036224.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Preliminary experiments gave an approx 13kDa band in Human Peripheral Blood Lymphocyte lysates after 0.01µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 22.6kDa according to NP_036224.1. The 13kDa band was successfully blocked by incubation with the immunizing peptide.

IHC: In paraffin embedded Human Lymph Node shows membrane staining just outside the mantle zone of the Germinal Centre. Recommended concentration: 1-2µg/ml. Paraffin embedded Human Tonsil and Thymus. Recommended concentration: 3.75µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human




EB09484 (2μg/ml) staining of paraffin embedded Human Lymph Node. Microwaved antigen retrieval with citrate buffer pH 6, HRP-staining.



EB09484 Negative Control showing staining of paraffin embedded Human Lymph Node, with no primary antibody.



EB09484 (3.75µg/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB09484 (3.75µg/ml) staining of paraffin embedded Human Thymus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.