



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12839 - Goat Anti-ICOS ligand / B7-H2 Antibody

Size: 100µg specific antibody in 200µl

Target Protein

Principal Names: ICOSLG, inducible T-cell co-stimulator ligand, B7-H2, B7H2, B7RP-1, B7RP1, CD275, GL50, ICOS-L, ICOSL, LICOS, B7 homolog 2, B7 homologue 2, B7-like protein GI50, B7-related protein 1, ICOS ligand, transmembrane protein B7-H2 ICOS ligand

Official Symbol: ICOSLG

Accession Number(s): NP_056074.1; NP_001269979.1; NP_001269980.1; NP_001269981.1

Human GeneID(s): [23308](#)

Important Comments: This antibody is expected to recognize all reported isoforms (NP_056074.1; NP_001269979.1; NP_001269980.1; NP_001269981.1).

Immunogen

Peptide with sequence C-SQTGNDIGERDKIT, from the internal region of the protein sequence according to NP_056074.1; NP_001269979.1; NP_001269980.1; NP_001269981.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Approx 33kDa band observed in Human Lymph Node lysates (calculated MW of 33.3kDa according to NP_056074.1). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB12839 (1 μ g/ml) staining of Human Lymph Node lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.