



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05586 - Goat Anti-ICSBP1 / IRF8 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** IRF8, ICSBP1, interferon consensus sequence binding protein 1, IRF-8, H-ICSBP, interferon regulatory factor 8

**Official Symbol:** IRF8

**Accession Number(s):** NP\_002154.1

**Human GeneID(s):** [3394](#)

**Non-Human GeneID(s):** 15900 (mouse)

### Immunogen

Peptide with sequence C-QRSFFRENQQITV, from the C Terminus of the protein sequence according to NP\_002154.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:8000.

**Western blot:** Approx 45-50kDa band seen in K562 lysate (Predicted MW of 47kDa according to NP\_002154.1). Recommended concentration: 0.5-2µg/ml.

**IHC:** Paraffin embedded Human Tonsil. Recommended concentration: 5µg/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Cow



EB05586 staining (1 $\mu$ g/ml) of K562 lysate (RIPA buffer, 30 $\mu$ g total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.



EB05586 (5 $\mu$ g/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.