

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB10997 - Goat Anti-IDH3B (aa369-383) Antibody

Size: 100µg specific antibody in 200µl



#### **Target Protein**

**Principal Names:** IDH3B, isocitrate dehydrogenase 3 (NAD+) beta, FLJ11043, H-IDHB, MGC903, RP46, NAD(+)-specific ICDH subunit beta, NAD+-specific ICDH, NAD+-specific isocitrate dehydrogenase b subunit, NAD+-specific isocitrate dehydrogenase beta, isocitrate dehydrogenase [NAD] subunit beta, mitochondrial, isocitrate dehydrogenase, NAD(+)-specific, mitochondrial, beta subunit, isocitric dehydrogenase subunit beta

Official Symbol: IDH3B

Accession Number(s): NP\_008830.2; NP\_777281.1

Human GenelD(s): 3420

Non-Human GenelD(s): 170718 (mouse), 94173 (rat)

Important Comments: This antibody is expected to recognize isoform a (NP\_008830.2)

and isoform c (NP\_777281.1).

#### **Immunogen**

Peptide with sequence C-TTDFIKSVIGHLQTK, from the C Terminus of the protein sequence according to NP\_008830.2; NP\_777281.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

**Western blot:** Approx 40kDa band observed in lysates of Mouse fetal Kidney and fetal Skeletal Muscle and of cell line NIH3T3 (calculated MW of 42.2kDa according to Human NP\_008830.2 and Mouse NP\_570954.1). Recommended concentration: 0.03-0.1µg/ml.

## **Species Reactivity**

Tested: Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB10997 ( $0.03\mu g/ml$ ) staining of Mouse fetal Kidney lysate ( $35\mu g$  protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.