

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona, CA 92065 USA

Inquiries: info@everestbiotech.com Sales: usasales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: 888-320-4628 (toll-free) Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07088 - Goat Anti-IFT88 / Polaris Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: IFT88, polaris, TTC10, tetratricopeptide repeat domain 10, HGNC:20606, D13S1056E, DAF19, IFT88, MGC26259, RP11-172H24.2, TG737, hTg737, TPR repeat protein 10, Probe hTg737 (polycystic kidney disease, autosomal recessive), Tg737 protein, intraflagellar transport 88 homolog, recessive polycystic kidney disease protein Tg737 homolog, intraflagellar transport 88 homolog (Chlamydomonas) **Official Symbol:** IFT88

Accession Number(s): NP_783195.2; NP_006522.2, NP_001340507.1; NP_001340498.1

Human GeneID(s): 8100

Non-Human GeneID(s): 21821 (mouse), 305918 (rat) Important Comments: This antibody is expected to recognise many reported isoforms.

Immunogen

Peptide with sequence C-KKRIDEDDFADEE, from the C Terminus of the protein sequence according to NP_783195.2; NP_006522.2, NP_001340507.1; NP_001340498.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 90kDa band observed in lysates of cell line A549 and in Mouse Testis lysates, and in preliminary testing of Rat Testes lysate (calculated MW of 90.4kDa according to Hunab NP_001340498.1, and 93.1kDa according to Mouse NP_033402.2 93.). Recommended concentration: 1-3ug/ml. Primary incubation 1 hour at room temperature.

Immunofluorescence: Strong expression of the protein seen in the nucleus and less staining in the cytoplasm of HepG2 cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of HepG2 cells. Recommended concentration: 10ug/ml.

Species Reactivity

Tested: Human, Mouse Expected from sequence similarity: Human, Mouse, Rat, Dog


Specific Reference


This antibody (previous batch) has been successfully used in IF on Mouse:


Miyoshi K, Kasahara K, Miyazaki I, Shimizu S, Taniguchi M, Matsuzaki S, Tohyama M, Asanuma M.

Pericentrin, a centrosomal protein related to microcephalic primordial dwarfism, is required for olfactory cilia assembly in mice.

FASEB J. 2009 Oct;23(10):3289-97. PMID: 19470799









EB07088 Immunofluorescence analysis of paraformaldehyde fixed HepG2 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG

(10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB07088 Flow cytometric analysis of paraformaldehyde fixed HepG2 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.



EB07088 (3µg/ml) staining of paraffin embedded Human Kidney. Microwaved antigen retrieval with Tris/EDTA buffer pH9, HRP-staining. **This data is from a previous batch, not on sale.**