

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08005 - Goat Anti-IGFBP3 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** IGFBP3, insulin-like growth factor binding protein 3, tcag7.703, BP-53, IBP3, IGF-binding protein 3, acid stable subunit of the 140 K IGF complex, binding protein 29, binding protein 53, growth hormone-dependent binding protein

Official Symbol: IGFBP3

Accession Number(s): NP\_001013416.1; NP\_000589.2

Human GeneID(s): 3486

Non-Human GenelD(s): 16009 (mouse), 24484 (rat)

Important Comments: This antibody is expected to recognise both reported isoforms

(NP\_001013416.1 and NP\_000589.2).

## **Immunogen**

Peptide with sequence C-RYKVDYESQSTDTQN, from the internal region of the protein sequence according to NP\_001013416.1; NP\_000589.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:64000.

**Western blot:** Approx 28kDa band observed in Human Breast Cancer lysates (calculated MW of 31.7kDa according to NP\_001013416.1 and 32.2kDa according to NP\_000589.2). Recommended concentration: 0.02-0.06µg/ml. An additional band of 50kDa was consistently observed, however this band was not blocked by the immunizing peptide and it is therefore a non-specific signal. We call for caution when used for other assays than Western blot.

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB08005 (0.02μg/ml) staining of Human Breast Cancer lysate (35μg protein in RIPA buffer) with (B) and without (A) blocking with the immunising peptide. Primary incubation was 1 hour. Detected by chemiluminescence.