

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB09753 - Goat Anti-INCA / CARD17 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: INCA, CARD17, caspase recruitment domain family, member 17, Inhibitory CARD, caspase recruitment domain-containing protein 17, caspase-1 inhibitor INCA, inhibitory caspase recruitment domain (CARD) protein, inhibitory caspase

recruitment domain protein
Official Symbol: INCA

Accession Number(s): NP_001007233.1

Human GeneID(s): 440068

Immunogen

Peptide with sequence C-NHLTTQDSQIVLPS, from the C Terminus of the protein sequence according to NP_001007233.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Approx 15kDa double band observed in Human Kidney and Lung lysates

(calculated MW of 11.9kDa according to NP_001007233.1). Recommended

concentration: 0.5-1.5µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB09753 (0.5μg/ml) staining of Human Kidney lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.