

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB08674-T - Goat Anti-IRAK3 (mouse) Antibody -

Size: 20µg specific antibody in 40µl



Principal Names: IRAK3, interleukin-1 receptor-associated kinase 3, ASRT5, IRAK-M,

IRAKM, interleukin-1 receptor-associated kinase M

Official Symbol: IRAK3

Accession Number(s): NP_082955.2

Non-Human GeneID(s): 73914 (mouse), 314870 (rat)

Immunogen

Peptide with sequence C-GHSYGSKPMEKR, from the internal region (near the C Terminus) of the protein sequence according to NP_082955.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 65-70kDa band observed in Rat Liver lysates, and in preliminary testing of Rat Adrenal Gland lysate (calculated MW of 69.0kDa according to Rat, NP_001101571.1 and 67.8kDa according to Mouse NP_082955.2). Recommended concentration: 0.03-0.1μg/ml. Primary incubation 1 hour at room temperature.

A customer detected a band at approx. 70kDa in Mouse Bone Marrow lysates, which was not present in the knock-out mouse.

Species Reactivity

Tested: Mouse, Rat

Expected from sequence similarity: Mouse, Rat

Specific Reference

This antibody has been successfully used in the following paper:

Gurcan Tunali, Marta Rúbies Bedós, Divya Nagarajan, Patrik Fridh, Irineos Papakyriacou, Yumeng Mao


IL-1 receptor-associated kinase-3 acts as an immune checkpoint in myeloid cells to limit cancer immunotherapy

J Clin Invest. 2023 Apr 3;133(7):e161084.

PMID: 36757800

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

B08674 (0.1ug/ml) staining of Rat Liver lysate (35µg protein in RIPA buffer). Detected with chemiluminescence.





 $EB08674\ (0.5\mu g/ml)\ staining\ of\ Mouse\ Bone\ Marrow\ (wildtype\ first\ lane,\ KO\ second\ lane)\ lysates\ (35\mu g\ protein\ lane)\ staining\ of\ Mouse\ Bone\ Marrow\ (wildtype\ first\ lane,\ kO\ second\ lane)\ lysates\ (35\mu g\ protein\ lysates\ ly$ in RIPA buffer). Primary incubation was overnight at 4C Detected with chemiluminescence. Data obtained from anonymous customer.