

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05585 - Goat Anti-IRF5 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

Principal Names: IRF5, interferon regulatory factor 5, SLEB10

Official Symbol: IRF5

Accession Number(s): NP\_116032.1; NP\_001092097.2; NP\_001092099.1; :

NP\_001229381.1

Human GeneID(s): 3663

 $\textbf{Important Comments:} \ \textbf{This antibody is expected to recognize reported isoforms}$ 

(NP\_116032.1; NP\_001092097.2; NP\_001092099.1; : NP\_001229381.1

#### **Immunogen**

Peptide with sequence C-QGPWPMHPAGMQ, from the C Terminus of the protein sequence according to NP\_116032.1; NP\_001092097.2; NP\_001092099.1; : NP\_001229381.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.


**Western blot:** Approx. 55+60kDa bands observed in Human Spleen lysates, approx. 65kDa in lysates of cell line A549, and approx. 70kDa in lysates of cell lines Daudi (calculated MW of 57.8kDa according to NP\_001092099.1). These molecular weights have been observed by other commercial sources, and were successfully blocked by incubation with the immunizing peptide. Recommended concentration: 1-2μg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Spleen. Recommended concentration: 5µg/ml.

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB05585 optimised QC. Primary incubation 1 hour at room temperature.

Images A+B: A549 cell lysate + peptide, incubation at primary Ab concentration 1ug/ml and C+D: Daudi cell lysate + peptide, incubation at primary Ab concentration 1ug/ml. (Loaded 35µg protein in RIPA buffer, per lane). Detected by chemiluminescence.




EB05585 optimised QC. Primary incubation 1 hour at room temperature.

Image A: Human Spleen lysate at primary Ab concentration 1ug/ml. (Loaded 35µg protein in RIPA buffer, per lane). Detected by chemiluminescence.



EB05585 ( $5\mu g/ml$ ) staining of paraffin embedded Human Spleen. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



 ${\tt EB05585\ Negative\ Control\ showing\ staining\ of\ paraffin\ embedded\ Human\ Spleen,\ with\ no\ primary\ antibody.}$