



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB12559 - Goat Anti-ITGB2 / LFA-1 (aa86-99) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** ITGB2, integrin, beta 2 (complement component 3 receptor 3 and 4 subunit), CD18, LAD, LCAMB, LFA-1, MAC-1, MF17, MFI7, cell surface adhesion glycoprotein LFA-1/CR3/P150,959 beta subunit precursor), cell surface adhesion glycoproteins LFA-1/CR3/p150,95 sub

**Official Symbol:** ITGB2

**Accession Number(s):** NP\_000202.2

**Human GeneID(s):** [3689](#)

**Important Comments:** Reported variants represent identical protein: NP\_001120963.1, NP\_000202.2.

### Immunogen

Peptide with sequence C-ETQEDHNGGQKQLS, from the internal region of the protein sequence according to NP\_000202.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:128000.


**Western blot:** Approx. 85kDa band observed in lysates of cell lines K562, MOLT4 and U937 (calculated MW of 84.8kDa according to NP\_000202.2). Recommended concentration: 0.5-2µg/ml. Primary incubation 1 hour at room temperature.

**Flow Cytometry:** Flow cytometric analysis of U937 cells. Recommended concentration: 10ug/ml.

### Species Reactivity


**Tested:** Human

**Expected from sequence similarity:** Human



EB12559 optimised QC. Primary incubation 1 hour at room temperature.

Image A: U937 cell lysate at primary Ab concentration 2ug/ml, Image B: K562 cell lysate at primary Ab concentration 0.5ug/ml and Image C: MOLT4 cell lysate at primary Ab concentration 1ug/ml. (Loaded 35µg protein in RIPA buffer, per lane). Detected by chemiluminescence.



EB12559 Flow cytometric analysis of paraformaldehyde fixed U937 cells (blue line) permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488® conjugated secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.