



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB08802 - Goat Anti-ITK Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ITK, IL2-inducible T-cell kinase, EMT, LYK, MGC126257, MGC126258, PSCTK2, homolog of mouse T-cell itk/tsk, tyrosine-protein kinase ITK/TSK, tyrosine-protein kinase LYK

Official Symbol: ITK

Accession Number(s): NP_005537.3

Human GeneID(s): [3702](#)

Immunogen

Peptide with sequence C-KERPEDRPAFSR, from the internal (near C terminus) of the protein sequence according to NP_005537.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 70kDa band observed in Rat Colon, Lung and Spleen lysates (calculated MW of 72.4kDa according to Rat NP_001102295.2). Recommended concentration: 0.3-1µg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Spleen. Recommended concentration: 3.75ug/ml.


Immunofluorescence: Strong expression of the protein seen in the cytoplasm and membranes of Jurkat cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of Jurkat cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Human, Rat


Expected from sequence similarity: Human, Rat, Cow




EB08802 (0.3 μ g/ml) staining of Rat Colon (A), (0.5 μ g/ml) Lung (B) and (1 μ g/ml) Spleen (C) lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB08802 (3.75 μ g/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB08802 Immunofluorescence analysis of paraformaldehyde fixed Jurkat cells immobilized on Shifix™ coverslip, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing membrane and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB08802 Flow cytometric analysis of paraformaldehyde fixed Jurkat cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.