

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

## EB08857 - Goat Anti-KCNMA1 Antibody

Size: 100µg specific antibody in 200µl



#### **Target Protein**

Principal Names: KCNMA1, potassium large conductance calcium-activated channel, subfamily M, alpha member 1, BKTM, DKFZp686K1437, KCa1.1, MGC71881, MaxiK, SAKCA, SLO, SLO-ALPHA, SLO1, mSLO1, BKCA alpha subunit, Drosophila slowpoke-like, OTTHUMP0000066198, OTTHUMP0000064154, OTTHUMP0000064155, OTTHUMP0000064156, OTTHUMP0000064157, OTTHUMP0000064158, OTTHUMP0000064159, OTTHUMP0000064160, OTTHUMP0000064161, OTTHUMP0000064162, OTTHUMP0000064164, OTTHUMP0000064165, bA205K10.1 (potassium large conductance calcium-activated channel, subfamily M, alpha member 1), large conductance calcium-activated potassium

Official Symbol: KCNMA1

Accession Number(s): NP\_001014797.1

Human GeneID(s): 3778

Non-Human GenelD(s): 16531 (mouse), 83731 (rat)

Important Comments: This antibody is expected to recognize isoform a

channel subfamily M alpha member 1, stretch-activated Kca channel

(NP\_001014797.1) only.

### **Immunogen**

Peptide with sequence C-RESRDKQNRKEMVYR, from the C Terminus of the protein sequence according to NP\_001014797.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:2000.

Western blot: Preliminary experiments in Human Brain (Cerebellum, Cerebral Cortex, Frontal Cortex) lysates gave no specific signal but low background (at antibody concentration up to 1μg/ml). We would appreciate any feedback from people in the field have any results been reported with other antibodies/lysates?

# **Species Reactivity**

Tested:

Expected from sequence similarity: Human, Mouse, Rat, Cow