

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB12963 - Goat Anti-Ketohexokinase Antibody

Size: 100µg specific antibody in 200µl



#### **Target Protein**

Principal Names: KHK, ketohexokinase (fructokinase), hepatic fructokinase,

ketohexokinase

Official Symbol: KHK

Accession Number(s): NP\_000212.1; NP\_006479.1

Human GeneID(s): 3795

Non-Human GenelD(s): 16548 (mouse), 25659 (rat)

Important Comments: This antibody is expected to recognize both reported isoforms

(NP\_000212.1; NP\_006479.1).

### Immunogen

Peptide with sequence RCLSQRWQRGGNASN, from the internal region of the protein sequence according to NP\_000212.1; NP\_006479.1.

Please note the peptide is available for sale.

#### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot:** Approx 30kDa band observed in Human, Mouse, Rat and Pig Liver lysates (calculated MW of 32.7kDa according to Human NP\_000212.1, Mouse NP\_032465.2, Rat NP\_114061.1 and of 32.6kDa according to Pig XP\_003125348.1). Recommended


concentration: 0.01-0.03µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Kidney. Recommended concentration: 5µg/ml.


#### **Species Reactivity**

Tested: Human, Mouse, Rat, Pig

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB12963 (0.03μg/ml) staining of Human (A), Mouse (B), Rat (C) and Pig (D) Liver lysates (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB12963 (5μg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.