



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06238 - Goat Anti-KPNA4 / IPOA3 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: KPNA4, IPOA3, QIP1, SRP3, MGC12217, MGC26703, karyopherin alpha 4 (importin alpha 3), importin alpha 3, importin-alpha-Q1, karyopherin alpha 4

Official Symbol: KPNA4

Accession Number(s): NP_002259.1

Human GeneID(s): [3840](#)

Immunogen

Peptide with sequence C-NSSANVPTEGFQF, from the C Terminus of the protein sequence according to NP_002259.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 55kDa band observed in Human and Mouse Ovary and in Mouse Testis lysates, and approx 60kDa in Rat Testis lysates (calculated MW of 57.9kDa according to Human NP_002259.1, Mouse NP_032493.1 and Rat NP_001014793.1). Recommended concentration: 0.03-0.3µg/ml. Primary incubation 1 hour at room temperature.

Species Reactivity

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

Specific References

This antibody (previous batch) has been successfully used in the following paper:

Sadequl Islam, Yang Sun, Yuan Gao, Tomohisa Nakamura, Arshad Ali Noorani, Tong Li, Philip C Wong, Noriyuki Kimura, Etsuro Matsubara, Kensaku Kasuga, Takeshi Ikeuchi, Taisuke Tomita, Kun Zou, Makoto Michikawa

Presenilin Is Essential for ApoE Secretion, a Novel Role of Presenilin Involved in Alzheimer's Disease Pathogenesis.

J Neurosci. 2022 Feb 23;42(8):1574-1586.

PMID: 34987110

This antibody (previous batch) has been successfully used in Western blot on Human and Mouse:

Huenniger K, Krämer A, Soom M, Chang I, Köhler M, Depping R, Kehlenbach RH, Kaether C.

Notch1 signaling is mediated by importins alpha 3, 4, and 7.

Cell Mol Life Sci. 2010 Sep;67(18):3187-96.

PMID: 20454918



EB06238 staining (0.1 $\mu\text{g/ml}$) of Human Ovary (A), Mouse Testes (B), (0.3 $\mu\text{g/ml}$) Mouse Ovary (C) and (0.03 $\mu\text{g/ml}$) Rat Testes (D) lysate (RIPA buffer, 35 μg total protein per lane). Detected by chemiluminescence.