

International Office

Everest Biotech Ltd Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com Technical Service: technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06418-T - Goat Anti-LAT1 Antibody - Trial

Size: 20µg specific antibody in 40µl

Target Protein

Principal Names: LAT1, LAT, linker for activation of T cells, pp36, 36 kDa phospho-tyrosine adaptor protein, linker for activation of T-cells family member 1 Official Symbol: LAT

Accession Number(s): NP_055202.1; NP_001014987.1; NP_001014989.1; NP_001014988.1

Human GeneID(s): 27040

Important Comments: This antibody is expected to recognize isoform a (NP_055202.1), isoform b (NP_001014987.1 and NP_001014989.1) and isoform c (NP_001014988.1).

Immunogen

Peptide with sequence C-GAPDYENLQELN, from the C Terminus of the protein sequence according to NP_055202.1; NP_001014987.1; NP_001014989.1; NP_001014988.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.


Applications Tested

Peptide ELISA: antibody detection limit dilution 1:2000.

Western blot: Approx 37kDa band observed in Human Peripheral Blood Mononucleocytes lysates (calculated MW of 27.9kDa according to NP_055202.1). This molecular weight is routinely observed by other sources. Recommended concentration: 0.01-0.03µg/ml.

Species Reactivity

Tested: Human Expected from sequence similarity: Human, Mouse, Rat





| | 250kDa 150kDa 100kDa 75kDa |
|---|-------------------------------------|
| | 50kDa |
| - | 37kDa |
| | 25kDa |
| | 20kDa |
| | 15kDa |

EB06418 (0.01µg/ml) staining of Peripheral Blood Mononucleocytes lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.