



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06762 - Goat Anti-Lipocalin 2 / NGAL Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: NGAL, lipocalin 2 (oncogene 24p3), HGNC:6526, LCN2, uterocalin, oncogenic lipocalin 24p3, neutrophil gelatinase-associated lipocalin, lipocalin 2, MSFI, OTTHUMP00000022240, migration-stimulating factor inhibitor, neutrophil gelatinase-associated lipocalin, oncogene 24p3, siderocalin

Official Symbol: LCN2

Accession Number(s): NP_005555.2

Human GeneID(s): [3934](#)

Immunogen

Peptide with sequence C-AILREDKDPQKMY, from the internal region of the protein sequence according to NP_005555.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 22kDa band observed in Human Kidney lysates (calculated MW of 22.6kDa according to NP_005555.2). Recommended concentration: 0.01-0.03µg/ml.

IHC: In paraffin embedded Human Spleen shows staining of cytoplasm in mostly binucleated cells, and of nucleoli in some of the non-dividing cells. Recommended concentration: 2-3µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB06762 (0.03 μ g/ml) staining of Human Kidney lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB06762 (2.5 μ g/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.