

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05439 - Goat Anti-LIS1 / PAFAH1B1 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** PAFAH1B1, LIS1, platelet-activating factor acetylhydrolase, isoform lb, alpha subunit 45kDa, MDCR, PAFAH, lissencephaly 1 protein, Platelet-activating factor acetylhydrolase, isoform 1B, alpha subunit, platelet-activating factor acetylhydrolase, isoform lb, alpha subunit (45kD), platelet-activating factor acetylhydrolase, isoform lb,

subunit 1 (45kDa), LIS2, MDS **Official Symbol:** PAFAH1B1

Accession Number(s): NP\_000421.1

Human GeneID(s): 5048

## Immunogen

Peptide with sequence TGSVDQTVKVWECR, from the C Terminus of the protein sequence according to NP\_000421.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.


**Western blot:** Approx 40kDa band observed in Human Lung and Brain (Cerebellum) and in Rat Ovary lysates (calculated MW of 46.6kDa according to Human NP\_000421.1 and to Rat NP\_113951.1). Recommended concentration: 0.3-1µg/ml.

**IHC:** Paraffin embedded Human Brain (Cortex), Thyroid and Pancreas. Recommended concentration: 3.75µg/ml.


## **Species Reactivity**

Tested: Human, Rat


Expected from sequence similarity: Human, Rat




EB05439 (0.3 $\mu$ g/ml) staining of Rat Ovary lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB05439 (3.75 $\mu$ g/ml) staining of paraffin embedded Human Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB05439 (3.75 $\mu$ g/ml) staining of paraffin embedded Human Thyroid. Steamed antigen retrieval with citrate buffer Ph 6, AP-staining.



EB05439 (3.75 $\mu$ g/ml) staining of paraffin embedded Human Pancreas. Steamed antigen retrieval with citrate buffer Ph 6, AP-staining.