



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06545 - Goat Anti-LRP6 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: LRP6, low density lipoprotein receptor-related protein 6, low density lipoprotein-related protein 6

Official Symbol: LRP6

Accession Number(s): NP_002327.2

Human GeneID(s): [4040](#)

Non-Human GeneID(s): 16974 (mouse)

Important Comments: This antibody is expected to recognise an epitope corresponding to aa 1546-1560 of human LRP6 protein.

Immunogen

Peptide with sequence CMTSVATAKGYTSDL, from the internal region of the protein sequence according to NP_002327.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.


IHC: In paraffin embedded Human Kidney shows luminal membrane staining in DCT.
Recommended concentration: 4-6µg/ml.

Flow Cytometry: Flow cytometric analysis of K562 cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB06545 (4 μ g/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, HRP-staining.



EB06545 Flow cytometric analysis of paraformaldehyde fixed K562 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (1 μ g/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.