



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB08228 - Goat Anti-MAN2A1 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** MAN2A1, mannosidase, alpha, class 2A, member 1, GOLIM7, MANA2, MANII, Golgi alpha-mannosidase II, golgi integral membrane protein 7, mannosidase, alpha type II, mannosyl-oligosaccharide 1,3-1,6-alpha-mannosidase

**Official Symbol:** MAN2A1

**Accession Number(s):** NP\_002363.2

**Human GeneID(s):** [4124](#)

### Immunogen

Peptide with sequence C-KISSDIKSQNR, from the internal region of the protein sequence according to NP\_002363.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx. 150kDa band observed in Human Placenta lysates (calculated MW of 131kDa according to NP\_002363.2). In transfected HEK293 transiently expressing MAN2A1 a band of approx. 98kDa is observed. This band is not observed in the non-transfected HEK293. Recommended concentration: 1-3µg/ml. An anonymous customer found positive results in WB on Human dermal fibroblasts.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB08228 (0.3µg/ml) staining of Human Placenta lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



HEK293 overexpressing Man2A1 (RC220186) and probed with EB08228 (mock transfection in first lane), tested by Origene.