

#### International Office

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05965 - Goat Anti-MDA5 / IFIH1 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** IFIH1, MDA5, MDA-5, Hlcd, MGC133047, IDDM19, melanoma differentiation associated protein-5, DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide,

interferon induced with helicase C domain 1, helicard

Official Symbol: IFIH1

Accession Number(s): NP\_071451.2

Human GeneID(s): 64135

### **Immunogen**

Peptide with sequence SNGYSTDENFRYL-C, from the N Terminus of the protein sequence according to NP\_071451.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: This antibody has been successfully used in Western blot on African Green

Monkey: Sato S et al, Arthritis Rheum. 2009 Jul;60(7):2193-200. PMID: 19565506.

IHC: In paraffin embedded Human Pancreas, shows strong cytoplasmic staining of some

cells in the islets of Langerhans. Recommended concentration: 10ug/ml.

**Flow Cytometry:** Flow cytometric analysis of K562 cells. Recommended concentration: 10ug/ml.

**Species Reactivity** 

Tested: Human, Monkey

Expected from sequence similarity: Human


### **Specific Reference**

This antibody has been successfully used in Western blot on African Green Monkey:


Sato S, Hoshino K, Satoh T, Fujita T, Kawakami Y, Fujita T, Kuwana M. RNA helicase encoded by melanoma differentiation-associated gene 5 is a major autoantigen in patients with clinically amyopathic dermatomyositis: Association with rapidly progressive interstitial lung disease.

Arthritis Rheum. 2009 Jul;60(7):2193-200.

PMID: 19565506



EB05965 (10µg/ml) staining of paraffin embedded Human Pancreas. Microwaved antigen retrieval with Tris/EDTA buffer pH9, HRP-staining.



EB05965 Flow cytometric analysis of paraformaldehyde fixed K562 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.