

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06765 - Goat Anti-MIF Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** MIF, macrophage migration inhibitory factor (glycosylation-inhibiting factor), HGNC:7097, GIF, GLIF, MMIF, glycosylation-inhibiting factor, phenylpyruvate tautomerase, L-dopachrome isomerase, L-dopachrome tautomerase

Official Symbol: MIF

Accession Number(s): NP\_002406.1

Human GeneID(s): 4282

#### **Immunogen**

Peptide with sequence C-NAANVGWNNSTFA, from the C Terminus of the protein sequence according to NP\_002406.1.

Please note the <u>peptide</u> is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.


Western blot: Approx 13kDa band observed in Human Lymph Node, and Thymus lysates and similar bands are observed in lysates of cell lines Daudi and Jurkat (calculated MW of 12.5kDa according to NP\_002406.1). Recommended concentration: 0.01-0.03μg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Thymus. Recommended concentration: 5µg/ml.


#### **Species Reactivity**


Tested: Human

Expected from sequence similarity: Human, Rat, Mouse



EB06765 ( $0.01\mu g/ml$ ) staining of Human Thymus lysate ( $35\mu g$  protein in RIPA buffer). Detected by chemiluminescence.





EB06765 ( $5\mu g/ml$ ) staining of paraffin embedded Human Thymus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.