

### International Office

### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB11928 - Goat Anti-MK5 / MAPKAPK5 (aa192-206) Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

Principal Names: MAPKAPK5, mitogen-activated protein kinase-activated protein kinase

5, MAPKAP-K5, MK-5, MK5, PRAK, MAP kinase-activated protein kinase 5,

MAPK-activated protein kinase 5, MAPKAP kinase 5, MAPKAPK-5,

p38-regulated/activated protein kinase

Official Symbol: Mapkapk5

Accession Number(s): NP\_003659.2; NP\_620777.1

Human GenelD(s): 8550

Non-Human GeneID(s): 17165 (mouse), 498183 (rat)

## **Immunogen**

Peptide with sequence C-PQVLEAQRRHQKEKS, from the internal region of the protein sequence according to NP\_003659.2; NP\_620777.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:64000.

**Western blot:** In transfected HEK293 transiently expressing either Mouse MK5 fused to GFP or the similar Mouse MK2 fused to GFP, only the expected band was observed in the MK5 lane after EB11928 labeling. Data obtained from Dr. M. B. Menon, Inst. Biochemistry, Hannover Medical School, Germany. Recommended concentration, 0.5-1µg/ml.


**IHC:** In paraffin embedded Human Kidney shows nuclear staining in the DCT. Recommended concentration, 4-8µg/ml.

Immunoprecipitation: MK5/ERK3 double knockout Mouse Embryonic Fibroblasts (MEFs) retrovirally transduced wiith MK5/ERK3 or empty vector (GFP) were lysed from confluent plates and used for IP with 1.5ug EB11927 or EB11928. Western blots of the IP were labelled with mouse anti-MK5 (SC) or with rabbit anti-ERK3 serum (CST). An approx 55kDa of MK5 is only precipitated from lysates of those KO MEFs that have been rescued by the ERK3/MK5 expression construct as described in PMID: 22508986. Data obtained from Dr. M. B. Menon, Inst. Biochemistry, Hannover Medical School, Germany.


# **Species Reactivity**

Tested: Human, Mouse


Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



HEK293 lysate (10ug protein in RIPA buffer) overexpressing Mouse MK5-GFP (first lane) or Mouse MK2-GFP (second lane) probed with EB11927 (0.5ug/ml) in right panel and with EB11928 (0.5ug/ml) on left panel, Primary incubations were for 2 hours. Detected by chemiluminescence.



EB11927 and EB11928 (1.5ug) immunoprecipitations from lysates of MK5/ERK3 double knockout MEFs, with (third and fifth lanes) and without (fourth and sixth lanes) rescued MK5/ERK3 expression through retroviral transduction. The corresponding lysates (first and second lane resp.) were analyzed in parallel in this Western blot labelled with mouse anti-MK5 / PRAK (and co-precipitation was measured using rabbit anti-ERK3 in the lower panel).



EB11927 (5µg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.