

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05530 - Goat Anti-MLH3 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** MLH3, mutL homolog 3 (E. coli), HNPCC, HNPCC7, MGC138372, S240II117, mutL (E. coli) homolog 3, mismatch repair gene MLH3, DNA mismatch repair

protein MLH3, mutL homolog 3

Official Symbol: MLH3

Accession Number(s): NP\_001035197.1; NP\_055196.2

Human GeneID(s): 27030

Important Comments: This antibody is expected to recognize isoform 1

(NP\_001035197.1) and isoform 2 (NP\_055196.2).

## Immunogen

Peptide with sequence RQSLQQSMPPCEPP, from the C Terminus of the protein sequence according to NP\_001035197.1; NP\_055196.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:2000.


**Immunofluorescence:** Strong expression of the protein seen in the nuclei of HeLa cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of HeLa cells. Recommended concentration: 10ug/ml.


#### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB05530 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear staining. Actin filaments were stained with phalloidin (red) and the nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB05530 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control:

Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.