

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05729 - Goat Anti-KMT2B/MLL4 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** KMT2B, MLL4, HRX2, MLL2, WBP7, DYT28, MLL1B, myeloid/lymphoid or mixed-lineage leukemia 4, CXXC10, histone-lysine

N-methyltransferase 2B, KIAA0304, Lysine N-methyltransferase 2B, mixed lineage

leukemia 4, WBP-7, WBP-7 **Official Symbol:** KMT2B

Accession Number(s): NP\_055542.1

Human GeneID(s): 9757

Non-Human GenelD(s): 75410 (mouse)

#### **Immunogen**

Peptide with sequence CNCGAKRCRRFLN, from the C Terminus of the protein sequence according to NP\_055542.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.


**Western blot:** Using our routine western blotting protocol we would not expect to reliably detect proteins larger than 250kDa (calculated MWt. of approx. 294KDa according to NP\_055542.1).

**IHC:** Positive staining seen in paraffin embedded Human Testis and Human Brain Cortex. Recommended concentration: 5-10µg/ml.


## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig



EB05729 (5μg/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB05729 (5μg/ml) staining of paraffin embedded Human Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.