



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11719 - Goat Anti-MMP3 (aa459-473) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CHDS6, matrix metallopeptidase 3 (stromelysin 1, progelatinase), matrix metalloproteinase 3 (stromelysin 1, progelatinase), matrix metalloproteinase-3, MGC126102, MGC126103, MGC126104, MMP3, MMP-3, OTTHUMP00000045949, proteoglycan, SL-1, STMY, STMY1, STR1

Official Symbol: MMP3

Accession Number(s): NP_002413.1

Human GeneID(s): [4314](#)

Immunogen

Peptide with sequence C-DPNAKKVTHLKSNS, from the C Terminus of the protein sequence according to NP_002413.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.


Western blot: Approx 55kDa band observed in Human Prostate and approx. 60kDa band in Mouse Eye lysates (calculated MW of 54.0 kDa according to Human NP_002413.1 and Mouse NP_034939.1). Recommended concentration: 0.1-0.3µg/ml. An additional band of unknown identity was also consistently observed at 24kDa in Human. This band was successfully blocked by incubation with the immunizing peptide.

IHC: Paraffin embedded Human Kidney. Recommended concentration: 4-6µg/ml.


Species Reactivity

Tested: Human, Mouse


Expected from sequence similarity: Human




EB11719 (0.2 μ g/ml) staining of Human Prostate lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB11719 (0.3 μ g/ml) staining of Mouse Eye lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB11719 (6µg/ml) staining of paraffin embedded Human Kidney. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB11719 Negative Control showing staining of paraffin embedded Human Kidney, with no primary antibody.