

UK Office

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: <u>sales@everestbiotech.com</u> Tech support: <u>support@everestbiotech.com</u>

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore 405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries: info@everestbiotech.com Sales: usasales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: 888-320-4628 (toll-free) Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB10972 - Goat Anti-MNSOD (aa119-130) Antibody 🏹

Size: 100µg specific antibody in 200µl

Target Protein

Principal Names: SOD2, superoxide dismutase 2, mitochondrial, RP1-56L9.2, IPOB, MNSOD, MVCD6, Mn superoxide dismutase, indophenoloxidase B,

manganese-containing superoxide dismutase, superoxide dismutase [Mn], mitochondrial **Official Symbol:** SOD2

Accession Number(s): NP_000627.2; NP_001019637.1

Human GeneID(s): 6648

Non-Human GeneID(s): 20656 (mouse), 24787 (rat)

Important Comments: NB: The immunizing peptide represents the acetylation site including K122 according to isoform A. This antibody is expected to recognize both reported isoforms (NP_000627.2; NP_001019637.1) . Reported variants represent identical protein: NP_000627.2; NP_001019637.1.

Immunogen

Peptide with sequence C-EAIKRDFGSFDK, from the internal region of the protein sequence according to NP_000627.2; NP_001019637.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.


Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.


Western blot: Approx 24kDa band observed in Human Brain (cerebellum) lysates and in Mouse and Rat Brain and Spinal Cord lysates, while approx 26kDa band was observed in lysates of cell lines HeLa, HepG2, HEK293 and NIH3T3 (calculated MW of 24.8kDa according to Human NP_000627.2, and 24.6kDa according to Mouse NP_038699.2). Recommended concentration: 0.01-0.1µg/ml. Primary incubation was 1 hour. **IHC:** Paraffin embedded Human Skeletal Muscle. Recommended concentration: 5µg/ml.

Species Reactivity


Tested: Human, Mouse, Rat, Pig Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow, Zebrafish




EB10972 (0.3µg/ml) staining of Human Cerebellum (A), Mouse Brain (B), Rat Brain (C) and Pig Brain (D) lysates (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB10972 (0.1µg/ml) staining of HeLa, HepG2, HEK293 and NIH3T3 (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB10972 (0.01µg/ml) staining of Mouse (A) and Rat (B) Spinal Cord lysates (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB10972 (5µg/ml) staining of paraffin embedded Human Skeletal Muscle. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.